

Preprint, arXiv:1011.3487

SOME NEW SUPER-CONGRUENCES MODULO PRIME POWERS

ZHI-WEI SUN

Department of Mathematics, Nanjing University
Nanjing 210093, People's Republic of China
zwsun@nju.edu.cn
<http://math.nju.edu.cn/~zwsun>

ABSTRACT. Let $p > 3$ be a prime. We show that

$$\sum_{k=0}^{p-1} \binom{-1/(p+1)}{k}^{p+1} \equiv 0 \pmod{p^5} \quad \text{and} \quad \sum_{k=0}^{p-1} \binom{1/(p-1)}{k}^{p-1} \equiv 0 \pmod{p^4}.$$

For any positive integer $m \not\equiv 0 \pmod{p}$, we prove that

$$\sum_{k=0}^{p-1} (-1)^{km} \binom{p/m-1}{k}^m \equiv 0 \pmod{p^4},$$

and

$$\sum_{k=1}^{p-1} \frac{(-1)^{km}}{k^2} \binom{p/m-1}{k}^m \equiv \frac{1}{p} \sum_{k=1}^{p-1} \frac{1}{k} \pmod{p^3} \quad \text{if } p > 5.$$

The paper also contains some open conjectures.

1. INTRODUCTION

A p -adic congruence (with p a prime) is called a *super-congruence* if it happens to hold modulo higher powers of p . Here is a classical example due to J. Wolstenholme (cf. [W] or [HT]):

$$\sum_{k=1}^{p-1} \frac{1}{k} \equiv 0 \pmod{p^2} \quad \text{and} \quad \binom{2p-1}{p-1} \equiv 1 \pmod{p^3}$$

2010 *Mathematics Subject Classification*. Primary 11B65; Secondary 05A10, 11A07, 11B68.

Keywords. Super-congruences, binomial coefficients, Bernoulli numbers.

Supported by the National Natural Science Foundation (grant 10871087) and the Overseas Cooperation Fund (grant 10928101) of China.

for any prime $p > 3$. The reader may consult Long [L], Sun [Su1] and [Su2] for some known super-congruences.

In this paper we obtain some new super-congruences modulo prime powers motivated by the well-known formula

$$\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e.$$

Now we state our main results.

Theorem 1.1. *Let $p > 3$ be a prime. Then*

$$\sum_{k=0}^{p-1} \binom{-1/(p+1)}{k}^{p+1} \equiv 0 \pmod{p^5}. \quad (1.1)$$

Theorem 1.2. *Let $p > 3$ be a prime and let m be a positive integer not divisible by p . Then we have*

$$\sum_{k=0}^{p-1} (-1)^{km} \binom{p/m - 1}{k}^m \equiv 0 \pmod{p^4}. \quad (1.2)$$

In particular,

$$\sum_{k=0}^{p-1} \binom{1/(p-1)}{k}^{p-1} \equiv 0 \pmod{p^4}. \quad (1.3)$$

Remark 1.1. We conjecture that there are no composite numbers p satisfying (1.1) or (1.3). We also note that (1.1) and (1.3) can be refined as follows:

$$\sum_{k=0}^{p-1} \binom{-1/(p+1)}{k}^{p+1} \equiv \frac{p^5}{18} B_{p-3} \pmod{p^6} \quad \text{for any prime } p > 5, \quad (1.4)$$

and

$$\sum_{k=0}^{p-1} \binom{1/(p-1)}{k}^{p-1} \equiv \frac{2}{3} p^4 B_{p-3} \pmod{p^5} \quad \text{for any prime } p > 3, \quad (1.5)$$

where $B_0, B_1, B_2, \dots$ are Bernoulli numbers (see [IR, pp. 228–241] for an introduction to Bernoulli numbers). However, the proofs of (1.4) and (1.5) are too complicated.

Theorem 1.3. *Let $p > 3$ be a prime and let m be a positive integer not divisible by p .*

(i) *If $p > 5$, then*

$$\sum_{k=1}^{p-1} \frac{(-1)^{km}}{k^2} \binom{p/m-1}{k}^m \equiv \frac{1}{p} \sum_{k=1}^{p-1} \frac{1}{k} \pmod{p^3}. \quad (1.6)$$

Also, for any $n = 1, \dots, (p-3)/2$ we have

$$\sum_{k=1}^{p-1} \frac{(-1)^{km}}{k^{2n}} \binom{p/m-1}{k}^m \equiv -\frac{p}{2n+1} B_{p-1-2n} \pmod{p^2}. \quad (1.7)$$

(ii) *For every $n = 1, \dots, (p-3)/2$, we have*

$$\sum_{k=1}^{p-1} \frac{(-1)^{km}}{k^{2n-1}} \binom{p/m-1}{k}^m \equiv \left(1 + \frac{1-m}{2m}(2n+1)\right) \frac{p^2 n}{2n+1} B_{p-1-2n} \pmod{p^3}. \quad (1.8)$$

Remark 1.2. If n is a positive integer and $p > 2n+1$ is a prime then (1.8) with $m = p \pm 1$ yields

$$\sum_{k=1}^{p-1} \frac{1}{k^{2n-1}} \binom{1/(p-1)}{k}^{p-1} \equiv -\frac{2p^2 n^2}{2n+1} B_{p-1-2n} \pmod{p^3} \quad (1.9)$$

and

$$\sum_{k=1}^{p-1} \frac{1}{k^{2n-1}} \binom{-1/(p+1)}{k}^{p+1} \equiv \frac{p^2 n}{2n+1} B_{p-1-2n} \pmod{p^3}. \quad (1.10)$$

For a prime p and a p -adic number x , as usual we let $\nu_p(x)$ denote the p -adic valuation (i.e., p -adic order) of x .

Conjecture 1.1. *Let p be a prime and let n be a positive integer. Then*

$$\nu_p \left(\sum_{k=0}^{n-1} \binom{-1/(p+1)}{k}^{p+1} \right) \geq c_p \left\lfloor \frac{\nu_p(n) + 1}{2} \right\rfloor,$$

where

$$c_p = \begin{cases} 1 & \text{if } p = 2, \\ 3 & \text{if } p = 3, \\ 5 & \text{if } p \geq 5. \end{cases}$$

If $p > 3$ then

$$\nu_p \left(\sum_{k=0}^{n-1} \binom{1/(p-1)}{k}^{p-1} \right) \geq 4 \left\lfloor \frac{\nu_p(n) + 1}{2} \right\rfloor.$$

Now we raise one more conjecture.

Conjecture 1.2. *Let $m \geq 2$ and r be integers. And let $p > r$ be an odd prime not dividing m .*

(i) *If $m > 2$, $m \not\equiv r \pmod{2}$, and $p \equiv r \pmod{m}$ with $r \geq -m/2$, then*

$$\sum_{k=0}^{p-1} (-1)^{km} \binom{r/m}{k}^m \equiv 0 \pmod{p^3}. \quad (1.11)$$

(ii) *If $p \equiv r \pmod{2m}$ with $r \geq -m$, then*

$$\sum_{k=0}^{p-1} (-1)^k \binom{r/m}{k}^{2n+1} \equiv 0 \pmod{p^2} \quad \text{for all } n = 1, \dots, m-1. \quad (1.12)$$

Remark 1.3. The congruences in (1.11) and (1.12) modulo p are easy.

Theorems 1.1-1.2 and Theorem 1.3 will be proved in Sections 2 and 3 respectively.

2. PROOFS OF THEOREMS 1.1 AND 1.2

For $m = 1, 2, 3, \dots$ and $n = 0, 1, 2, \dots$, we define

$$H_n^{(m)} := \sum_{0 < k \leq n} \frac{1}{k^m}$$

and call it a harmonic number of order m . Those $H_n = H_n^{(1)}$ ($n = 0, 1, 2, \dots$) are usually called *harmonic numbers*.

Lemma 2.1. *Let $p > 3$ be a prime. Then*

$$H_{p-1} \equiv -\frac{p^2}{3} B_{p-3} \pmod{p^3}, \quad H_{p-1}^{(2)} \equiv \frac{2}{3} p B_{p-3} \pmod{p^2}, \quad (2.1)$$

and

$$\sum_{1 \leq i_1 < i_2 < \dots < i_k \leq p-1} \frac{1}{i_1 i_2 \dots i_k} \equiv \frac{(-1)^{k-1}}{k+1} p B_{p-1-k} \pmod{p^2} \quad (2.2)$$

for all $k = 1, 2, \dots, p-1$.

Remark 2.1. The results in Lemma 2.1 are known, see Theorem 5.1, Corollary 5.1 and the proof of [S, Theorem 6.1].

Lemma 2.2. *Let $p > 3$ be a prime. Then*

$$\sum_{k=1}^{p-1} H_k \equiv -\frac{p^3}{3} B_{p-3} - p + 1 \pmod{p^4}, \quad (2.3)$$

and

$$\sum_{k=1}^{p-1} H_k^{(2)} \equiv 0 \pmod{p^2} \quad \text{and} \quad \sum_{k=1}^{p-1} H_k^{(3)} \equiv 0 \pmod{p}. \quad (2.4)$$

Proof. For any positive integer m we have

$$\sum_{k=1}^{p-1} H_k^{(m)} = \sum_{k=1}^{p-1} \sum_{j=1}^k \frac{1}{j^m} = \sum_{j=1}^{p-1} \frac{\sum_{k=j}^{p-1} 1}{j^m} = \sum_{j=1}^{p-1} \frac{p-j}{j^m}.$$

Thus

$$\sum_{k=1}^{p-1} H_k = pH_{p-1} - p + 1, \quad \sum_{k=1}^{p-1} H_k^{(2)} = pH_{p-1}^{(2)} - H_{p-1}, \quad \sum_{k=1}^{p-1} H_k^{(3)} = pH_{p-1}^{(3)} - H_{p-1}^{(2)}.$$

Combining these with (2.1), we immediately get (2.3) and (2.4). $\square$

Lemma 2.3. *Let $p > 3$ be a prime. Then*

$$\sum_{k=1}^{p-1} \sum_{1 \leq i < j \leq k} \frac{1}{ij} \equiv -\frac{2}{3} p^2 B_{p-3} + p - 1 \pmod{p^3}, \quad (2.5)$$

$$\sum_{k=1}^{p-1} \sum_{1 \leq i_1 < i_2 < i_3 \leq k} \frac{1}{i_1 i_2 i_3} \equiv -\frac{p}{3} B_{p-3} - p + 1 \pmod{p^2}, \quad (2.6)$$

and

$$\sum_{k=1}^{p-1} \sum_{1 \leq i_1 < i_2 < i_3 < i_4 \leq k} \frac{1}{i_1 i_2 i_3 i_4} \equiv -1 \pmod{p}. \quad (2.7)$$

We also have

$$\sum_{k=1}^{p-1} \sum_{1 \leq i < j \leq k} \left(\frac{1}{ij^2} + \frac{1}{i^2 j} \right) \equiv 0 \pmod{p}. \quad (2.8)$$

Proof. For $s = 2, \dots, p-1$ it is clear that

$$\begin{aligned}
& \sum_{k=1}^{p-1} \sum_{1 \leq i_1 < i_2 < \dots < i_s \leq k} \frac{1}{i_1 i_2 \dots i_s} \\
&= \sum_{1 \leq i_1 < i_2 < \dots < i_s \leq p-1} \frac{\sum_{k=i_s}^{p-1} 1}{i_1 i_2 \dots i_s} = \sum_{1 \leq i_1 < i_2 < \dots < i_s \leq p-1} \frac{p - i_s}{i_1 i_2 \dots i_s} \\
&= {}^p \sum_{1 \leq i_1 < i_2 < \dots < i_s \leq p-1} \frac{1}{i_1 i_2 \dots i_s} - \sum_{1 \leq i_1 < i_2 < \dots < i_{s-1} \leq p-1} \frac{\sum_{i_{s-1} < i_s < p} 1}{i_1 i_2 \dots i_{s-1}} \\
&= {}^p \sum_{1 \leq i_1 < i_2 < \dots < i_s \leq p-1} \frac{1}{i_1 i_2 \dots i_s} - \sum_{1 \leq i_1 < i_2 < \dots < i_{s-1} \leq p-1} \frac{p - 1 - i_{s-1}}{i_1 i_2 \dots i_{s-1}}.
\end{aligned}$$

Thus, with the help of Lemmas 2.1, we have

$$\begin{aligned}
\sum_{k=1}^{p-1} \sum_{1 \leq i < j \leq k} \frac{1}{ij} &= {}^p \sum_{1 \leq i < j \leq p-1} \frac{1}{ij} - \sum_{i=1}^{p-1} \frac{p-1-i}{i} \\
&\equiv -\frac{p^2}{3} B_{p-3} - (p-1) H_{p-1} + p-1 \\
&\equiv -\frac{2}{3} p^2 B_{p-3} + p-1 \pmod{p^3}.
\end{aligned}$$

Also,

$$\begin{aligned}
& \sum_{k=1}^{p-1} \sum_{1 \leq i_1 < i_2 < i_3 \leq k} \frac{1}{i_1 i_2 i_3} \\
&= {}^p \sum_{1 \leq i_1 < i_2 < i_3 \leq p-1} \frac{1}{i_1 i_2 i_3} - \sum_{1 \leq i_1 < i_2 \leq p-1} \frac{p-1-i_2}{i_1 i_2} \\
&\equiv -(p-1) \sum_{1 \leq i_1 < i_2 \leq p-1} \frac{1}{i_1 i_2} + \sum_{i_1=1}^{p-1} \frac{\sum_{i_1 < i_2 < p} 1}{i_1} \\
&\equiv (p-1) \frac{p}{3} B_{p-3} + \sum_{i=1}^{p-1} \frac{p-1-i}{i} \equiv -\frac{p}{3} B_{p-3} - p+1 \pmod{p^2}
\end{aligned}$$

and

$$\begin{aligned}
& \sum_{k=1}^{p-1} \sum_{1 \leq i_1 < i_2 < i_3 < i_4 \leq k} \frac{1}{i_1 i_2 i_3 i_4} \\
&= p \sum_{1 \leq i_1 < i_2 < i_3 < i_4 \leq p-1} \frac{1}{i_1 i_2 i_3 i_4} - \sum_{1 \leq i_1 < i_2 < i_3 \leq p-1} \frac{p-1-i_3}{i_1 i_2 i_3} \\
&\equiv \sum_{1 \leq i_1 < i_2 < i_3 \leq p-1} \frac{1+i_3}{i_1 i_2 i_3} \equiv \sum_{1 \leq i_1 < i_2 \leq p-1} \frac{\sum_{i_2 < i_3 < p} 1}{i_1 i_2} \\
&\equiv \sum_{1 \leq i < j \leq p-1} \frac{p-1-j}{ij} \equiv - \sum_{i=1}^{p-1} \frac{\sum_{i < j < p} 1}{i} = - \sum_{i=1}^{p-1} \frac{p-1-i}{i} \equiv -1 \pmod{p}.
\end{aligned}$$

Finally, we note that

$$\begin{aligned}
& \sum_{k=1}^{p-1} \sum_{1 \leq i < j \leq k} \left(\frac{1}{ij^2} + \frac{1}{i^2 j} \right) = \sum_{1 \leq i < j \leq p-1} \left(\frac{p-j}{ij^2} + \frac{p-j}{i^2 j} \right) \\
&\equiv - \sum_{1 \leq i < j \leq p-1} \left(\frac{1}{ij} + \frac{1}{i^2} \right) \equiv - \sum_{i=1}^{p-1} \frac{\sum_{i < j < p} 1}{i^2} \\
&= - \sum_{i=1}^{p-1} \frac{p-1-i}{i^2} = H_{p-1} - (p-1)H_{p-1}^{(2)} \equiv 0 \pmod{p}
\end{aligned}$$

with the help of (2.1).

So far we have proved (2.5)-(2.8). $\square$

Proof of Theorem 1.1. For each $k = 1, \dots, p-1$, clearly

$$\begin{aligned}
& \left(\frac{-1/(p+1)}{k} \right)^{p+1} = \left(\frac{p/(p+1) - 1}{k} \right)^{p+1} = \prod_{j=1}^k \left(1 - \frac{p}{(p+1)j} \right)^{p+1} \\
&\equiv \prod_{j=1}^k \left(1 - \frac{(p+1)p}{(p+1)j} + \frac{(p+1)p}{2} \cdot \frac{p^2}{(p+1)^2 j^2} - \frac{(p+1)p(p-1)}{3!} \cdot \frac{p^3}{(p+1)^3 j^3} \right) \\
&= \prod_{j=1}^k \left(1 - \frac{p}{j} + \frac{p^3}{2(p+1)j^2} - \frac{p^4(p-1)}{6(p+1)^2 j^2} \right) \\
&\equiv \prod_{j=1}^k \left(1 - \frac{p}{j} + \frac{p^3(1-p)}{2j^2} + \frac{p^4}{6j^3} \right) \pmod{p^5}
\end{aligned}$$

and hence

$$\begin{aligned}
\binom{-1/(p+1)}{k}^{p+1} &\equiv 1 - pH_k + \frac{p^3(1-p)}{2}H_k^{(2)} + \frac{p^4}{6}H_k^{(3)} + p^2 \sum_{1 \leq i < j \leq k} \frac{1}{ij} \\
&\quad - p^3 \sum_{1 \leq i_1 < i_2 < i_3 \leq k} \frac{1}{i_1 i_2 i_3} + p^4 \sum_{1 \leq i_1 < i_2 < i_3 < i_4 \leq k} \frac{1}{i_1 i_2 i_3 i_4} \\
&\quad - \frac{p^4}{2} \sum_{1 \leq i < j \leq k} \left(\frac{1}{ij^2} + \frac{1}{i^2 j} \right) \pmod{p^5}.
\end{aligned}$$

Thus, in view of Lemmas 2.1-2.3, we obtain

$$\begin{aligned}
&\sum_{k=1}^{p-1} \binom{-1/(p+1)}{k}^{p+1} \\
&\equiv p - 1 - p \left(-\frac{p^3}{3} B_{p-3} - p + 1 \right) + p^2 \left(-\frac{p}{3} B_{p-3} - p + 1 \right) \\
&\quad - p^3 \left(-\frac{p}{3} B_{p-3} - p + 1 \right) - p^4 \\
&\equiv -1 \pmod{p^5}
\end{aligned}$$

and hence (1.1) follows.

The proof of Theorem 1.1 is now complete. $\square$

Proof of Theorem 1.2. For each $k \in \{1, \dots, p-1\}$, we have

$$\begin{aligned}
(-1)^{km} \binom{p/m-1}{k}^m &= \prod_{j=1}^k \left(1 - \frac{p}{jm} \right)^m \\
&\equiv \prod_{j=1}^k \left(1 - \frac{pm}{jm} + \frac{m(m-1)}{2} \cdot \frac{p^2}{j^2 m^2} - \binom{m}{3} \frac{p^3}{j^3 m^3} \right) \\
&\equiv \prod_{j=1}^k \left(1 - \frac{p}{j} + \frac{m-1}{2m} \cdot \frac{p^2}{j^2} - \frac{(m-1)(m-2)}{6m^2} \cdot \frac{p^3}{j^3} \right) \\
&\equiv 1 - pH_k + \frac{m-1}{2m} p^2 H_k^{(2)} - \frac{(m-1)(m-2)}{6m^2} p^3 H_k^{(3)} \\
&\quad + p^2 \sum_{1 \leq i < j \leq k} \frac{1}{ij} - p^3 \sum_{1 \leq i_1 < i_2 < i_3 \leq k} \frac{1}{i_1 i_2 i_3} \\
&\quad - \frac{m-1}{2m} p^3 \sum_{1 \leq i < j \leq k} \left(\frac{1}{ij^2} + \frac{1}{i^2 j} \right) \pmod{p^4}.
\end{aligned}$$

Therefore, applying Lemmas 2.2 and 2.3 we get

$$\sum_{k=1}^{p-1} (-1)^{km} \binom{p/m-1}{k}^m \equiv p - 1 - p(-p+1) + p^2(p-1) - p^3 = -1 \pmod{p^4}.$$

This proves (1.2). Clearly (1.2) in the case $m = p - 1$ yields (1.3). This concludes the proof. $\square$

3. PROOF OF THEOREM 1.3

Lemma 3.1. *Let m and n be positive integers with $m \leq 2n + 1$, and let $p > 2n + 1$ be a prime. Then*

$$\sum_{k=1}^{p-1} \frac{H_k^{(m)}}{k^{2n+1-m}} \equiv \frac{(-1)^{m-1}}{2n+1} \binom{2n+1}{m} B_{p-1-2n} \pmod{p}. \quad (3.1)$$

When $m < 2n$ we have

$$\sum_{k=1}^{p-1} \frac{H_k^{(m)}}{k^{2n-m}} \equiv \frac{pB_{p-1-2n}}{2n+1} \left(n + (-1)^m \frac{n-m}{m+1} \binom{2n+1}{m} \right) \pmod{p^2}. \quad (3.2)$$

Proof. Since $\sum_{k=1}^{p-1} k^s \equiv 0 \pmod{p}$ for any integer $s \not\equiv 0 \pmod{p-1}$ (see, e.g., [IR, p. 235]), we have $\sum_{k=1}^{p-1} 1/k^{2n+1} \equiv 0 \pmod{p}$. Hence

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{H_k^{(m)}}{k^{2n+1-m}} &= \sum_{k=1}^{p-1} \frac{1}{k^{2n+1}} + \sum_{k=1}^{p-1} \frac{1}{k^{2n+1-m}} \sum_{j=0}^{p-1} j^{p-1-m} \\ &\equiv \sum_{k=1}^{p-1} \frac{1}{k^{2n+1-m}(p-m)} \sum_{i=0}^{p-1-m} \binom{p-m}{i} B_i k^{p-m-i} \\ &\equiv -\frac{1}{m} \sum_{i=0}^{p-1-m} \binom{p-m}{i} B_i \sum_{k=1}^{p-1} k^{p-1-2n-i} \\ &\equiv \frac{1}{m} \sum_{\substack{0 \leq i \leq p-1-m \\ p-1|2n+i}} \binom{p-m}{i} B_i = \frac{1}{m} \binom{p-m}{2n+1-m} B_{p-1-2n} \\ &\equiv \frac{1}{m} \binom{-m}{2n+1-m} B_{p-1-2n} = \frac{(-1)^{m-1}}{2n+1} \binom{2n+1}{m} B_{p-1-2n} \pmod{p}. \end{aligned}$$

This proves (3.1).

Now assume that $m < 2n$. As $m, 2n - m \in \{1, \dots, p-2\}$, we have

$$\sum_{j=1}^{p-1} \frac{1}{j^m} \equiv \sum_{k=1}^{p-1} \frac{1}{k^{2n-m}} \equiv 0 \pmod{p}.$$

It is known that

$$\sum_{k=0}^{p-1} \frac{1}{k^s} \equiv \frac{ps}{s+1} B_{p-1-s} \pmod{p^2} \quad \text{for each } s = 1, \dots, p-2. \quad (3.3)$$

(See, e.g., [S, Corollary 5.1].) Thus

$$\sum_{j=1}^{p-1} \frac{1}{j^m} \sum_{k=1}^{p-1} \frac{1}{k^{2n-m}} + \sum_{k=1}^{p-1} \frac{1}{k^{2n}} \equiv \sum_{k=1}^{p-1} \frac{1}{k^{2n}} \equiv \frac{2n}{2n+1} pB_{p-1-2n} \pmod{p^2}.$$

On the other hand,

$$\begin{aligned} & \sum_{j=1}^{p-1} \frac{1}{j^m} \sum_{k=1}^{p-1} \frac{1}{k^{2n-m}} + \sum_{k=1}^{p-1} \frac{1}{k^{2n}} - \sum_{1 \leq j \leq k \leq p-1} \frac{1}{j^m k^{2n-m}} \\ &= \sum_{1 \leq k \leq j \leq p-1} \frac{1}{j^m k^{2n-m}} = \sum_{1 \leq j \leq k \leq p-1} \frac{1}{(p-j)^m (p-k)^{2n-m}} \\ &= \sum_{1 \leq j \leq k \leq p-1} \frac{(p+j)^m (p+k)^{2n-m}}{(p^2-j^2)^m (p^2-k^2)^{2n-m}} \\ &\equiv \sum_{1 \leq j \leq k \leq p-1} \frac{(j^m + pmj^{m-1})(k^{2n-m} + p(2n-m)k^{2n-m-1})}{j^{2m} k^{2(2n-m)}} \\ &\equiv \sum_{1 \leq j \leq k \leq p-1} \left(\frac{1}{j^m k^{2n-m}} + \frac{pm}{j^{m+1} k^{2n-m}} + \frac{p(2n-m)}{j^m k^{2n-m+1}} \right) \pmod{p^2}. \end{aligned}$$

Therefore, with the help of (3.1) we have

$$\begin{aligned} & \frac{2n}{2n+1} pB_{p-1-2n} - 2 \sum_{k=1}^{p-1} \frac{H_k^{(m)}}{k^{2n-m}} \\ &\equiv pm \sum_{k=1}^{p-1} \frac{H_k^{(m+1)}}{k^{2n-m}} + p(2n-m) \sum_{k=1}^{p-1} \frac{H_k^{(m)}}{k^{2n-m+1}} \\ &\equiv pm \frac{(-1)^m}{2n+1} \binom{2n+1}{m+1} B_{p-1-2n} + p(2n-m) \frac{(-1)^{m-1}}{2n+1} \binom{2n+1}{m} B_{p-1-2n} \\ &= (-1)^m \frac{2(m-n)}{m+1} \binom{2n+1}{m} \frac{pB_{p-1-2n}}{2n+1} \pmod{p^2} \end{aligned}$$

and hence (3.2) holds. $\square$

Remark 3.1. By [ST, (5.4)], $\sum_{k=1}^{p-1} H_k/k^2 \equiv B_{p-3} \pmod{p}$ for any prime $p > 3$. By [M, (5)], $\sum_{k=1}^{p-1} H_k/k^3 \equiv -pB_{p-5}/10 \pmod{p^2}$ for any prime $p > 5$. Obviously these two results are particular cases of Lemma 3.1.

Lemma 3.2. *Let $p > 5$ be a prime. Then*

$$\sum_{k=1}^{p-1} \frac{1 - pH_k}{k^2} \equiv \frac{H_{p-1}}{p} \pmod{p^3}. \quad (3.4)$$

Proof. In view of Theorem 5.1(a) and Remark 5.1 of [S],

$$\frac{H_{p-1}^{(2)}}{2} \equiv p \left(\frac{B_{2p-4}}{2p-4} - 2 \frac{B_{p-3}}{p-3} \right) \equiv -\frac{H_{p-1}}{p} \pmod{p^3}$$

and $H_{p-1}^{(3)} \equiv 0 \pmod{p^2}$. Also,

$$\sum_{k=1}^{p-1} \frac{H_{k-1}}{k^2} = \sum_{1 \leq j < k \leq p-1} \frac{1}{jk^2} \equiv -3 \frac{H_{p-1}}{p^2} \pmod{p^2}$$

by [T, Theorem 2.3]. So we have

$$\sum_{k=1}^{p-1} \frac{1 - pH_k}{k^2} = H_{p-1}^{(2)} - pH_{p-1}^{(3)} - p \sum_{k=1}^{p-1} \frac{H_{k-1}}{k^2} \equiv \frac{H_{p-1}}{p} \pmod{p^3}.$$

This concludes the proof. $\square$

Proof of Theorem 1.3. Let $k \in \{1, \dots, p-1\}$. By the proof of Theorem 1.2,

$$(-1)^{km} \binom{p/m-1}{k}^m \equiv 1 - pH_k + \frac{m-1}{2m} p^2 H_k^{(2)} + p^2 \sum_{1 \leq i < j \leq k} \frac{1}{ij} \pmod{p^3}.$$

Thus, for any given $n \in \{1, \dots, p-3\}$ we have

$$\begin{aligned} & \sum_{k=1}^{p-1} \frac{(-1)^{km}}{k^n} \binom{p/m-1}{k}^m \\ & \equiv \sum_{k=1}^{p-1} \frac{1 - pH_k}{k^n} + \frac{m-1}{2m} p^2 \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k^n} + \frac{p^2}{2} \sum_{k=1}^{p-1} \frac{H_k^2 - H_k^{(2)}}{k^n} \pmod{p^3}. \end{aligned} \tag{3.5}$$

(i) Now suppose that $1 \leq n \leq (p-3)/2$. Then $\sum_{k=1}^{p-1} H_k^{(2)}/k^{2n} \equiv 0 \pmod{p}$ since

$$2 \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k^n} \equiv \sum_{k=1}^{p-1} \left(\frac{H_k^{(2)}}{k^{2n}} + \frac{H_{p-k}^{(2)}}{(p-k)^{2n}} \right) \equiv \sum_{k=1}^{p-1} \frac{1/k^2}{k^{2n}} \equiv 0 \pmod{p}.$$

Thus

$$\sum_{k=1}^{p-1} \frac{(-1)^{km}}{k^{2n}} \binom{p/m-1}{k}^m \equiv \sum_{k=1}^{p-1} \frac{1 - pH_k}{k^{2n}} \pmod{p^2} \tag{3.6}$$

and

$$\sum_{k=1}^{p-1} \frac{(-1)^{km}}{k^{2n}} \binom{p/m-1}{k}^m \equiv \sum_{k=1}^{p-1} \frac{1-pH_k}{k^{2n}} + \frac{p^2}{2} \sum_{k=1}^{p-1} \frac{H_k^2}{k^{2n}} \pmod{p^3}. \quad (3.7)$$

By (3.3) and Lemma 3.1,

$$\sum_{k=1}^{p-1} \frac{1-pH_k}{k^{2n}} \equiv \left(\frac{2n}{2n+1} - 1 \right) pB_{p-1-2n} = -\frac{pB_{p-1-2n}}{2n+1} \pmod{p^2}.$$

So (1.7) follows from (3.6).

When $p > 5$, (3.7) in the case $n = 1$, together with (3.3) and the congruence

$$\sum_{k=1}^{p-1} \frac{H_k^2}{k^2} \equiv 0 \pmod{p}$$

(cf. [Su3, (1.5)]), yields (1.6).

(ii) Fix $n \in \{1, \dots, (p-3)/2\}$. Substituting $2n-1$ for n in (3.5) we get

$$\begin{aligned} & \sum_{k=1}^{p-1} \frac{(-1)^{km}}{k^{2n-1}} \binom{p/m-1}{k}^m \\ & \equiv \sum_{k=1}^{p-1} \frac{1-pH_k}{k^{2n-1}} + \frac{m-1}{2m} p^2 \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k^{2n-1}} + \frac{p^2}{2} \sum_{k=1}^{p-1} \frac{H_k^2 - H_k^{(2)}}{k^{2n-1}} \pmod{p^3}. \end{aligned} \quad (3.8)$$

In view of a known result (cf. [S, Theorem 5.1(a)]),

$$\sum_{k=1}^{p-1} \frac{1}{k^{2n-1}} \equiv \frac{n-2n^2}{2n+1} p^2 B_{p-1-2n} \pmod{p^3}.$$

By Lemma 3.1,

$$\sum_{k=1}^{p-1} \frac{H_k}{k^{2n-1}} \equiv \frac{1+3n-2n^2}{2(2n+1)} pB_{p-1-2n} \pmod{p^2}$$

and

$$\sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k^{2n-1}} \equiv -nB_{p-1-2n} \pmod{p}.$$

Note also that

$$\sum_{k=1}^{p-1} \frac{H_k^2}{k^{2n-1}} = \sum_{k=1}^{p-1} \frac{H_{p-k}^2}{(p-k)^{2n-1}} \equiv -\sum_{k=1}^{p-1} \frac{(H_k - 1/k)^2}{k^{2n-1}} \pmod{p}$$

and hence

$$\sum_{k=1}^{p-1} \frac{H_k^2}{k^{2n-1}} \equiv \sum_{k=1}^{p-1} \frac{H_k}{k^{2n}} - \frac{1}{2} \sum_{k=1}^{p-1} \frac{1}{k^{2n+1}} \equiv \sum_{k=1}^{p-1} \frac{H_k}{k^{2n}} \equiv B_{p-1-2n} \pmod{p}.$$

Combining all these we obtain (1.8) from (3.8).

The proof of Theorem 1.3 is now complete. $\square$

REFERENCES

- [HT] C. Helou and G. Terjanian, On Wolstenholme's theorem and its converse, J. Number Theory **128**(2008) 475–499.
- [IR] K. Ireland and M. Rosen, A Classical Introduction to Modern Number Theory (Graduate texts in math.; 84), 2nd ed., Springer, New York, 1990.
- [L] L. Long, *Hypergeometric evaluation identities and supercongruences*, Pacific J. Math. **249** (2011), 405–418.
- [M] R. Meštrović, *A proof of a conjecture by Sun on congruence for harmonic numbers*, preprint, arXiv:1108.1171. <http://arxiv.org/abs/1108.1171>.
- [S] Z. H. Sun, *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Discrete Appl. Math. **105** (2000), 193–223.
- [Su1] Z. W. Sun, *Binomial coefficients, Catalan numbers and Lucas quotients*, Sci. China Math. **53** (2010), 2473–2488..
- [Su2] Z. W. Sun, *On congruences related to central binomial coefficients*, J. Number Theory **131** (2011), in press.
- [Su3] Z. W. Sun, *Arithmetic theory of harmonic numbers*, Proc. Amer. Math. Soc., in press. <http://arxiv.org/abs/0911.4433>.
- [ST] Z. W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Appl. Math. **45** (2010), 125–148.
- [T] R. Tauraso, *More congruences for central binomial coefficients*, J. Number Theory **130** (2010), 2639–2649.
- [W] J. Wolstenholme, *On certain properties of prime numbers*, Quart. J. Appl. Math. **5** (1862), 35–39.